



Training the Immune System to Fight Cancer

John K. Celebi, MBA
President & Chief Executive Officer

September 2021

NASDAQ: SNSE



Disclaimer



This presentation has been prepared by Sensei Biotherapeutics, Inc. (the "Company," "we," "us") and is made for informational purposes only. The information set forth herein does not purport to be complete or to contain all of the information you may desire. Statements contained herein are made as of the date of this presentation unless stated otherwise, and neither the delivery of this presentation at any time, nor any sale of securities, shall under any circumstances create an implication that the information contained herein is correct as of any time after such date or that information will be updated or revised to reflect information that subsequently becomes available or changes occurring after the date hereof.

This presentation contains estimates and other statistical data made by independent parties and by us relating to market shares and other data about our industry. This presentation also contains "forward-looking" statements as that term is defined in the Private Securities Litigation Reform Act of 1995 that are based on our management's beliefs and assumptions and on information currently available to management. These forward-looking statements include, without limitation, statements regarding our industry, business strategy, plans, goals and expectations concerning our market position, product expansion, future operations, margins, profitability, future efficiencies, and other financial and operating information. When used in this presentation, the words "may," "believes," "intends," "seeks," "anticipates," "plans," "estimates," "expects," "should," "assumes," "continues," "could," "will," "future" and the negative of these or similar terms and phrases are intended to identify forward-looking statements. Forward-looking statements involve known and unknown risks, uncertainties and other factors that may cause our actual results, performance or achievements to be materially different from any future results, performance or achievements expressed or implied by the forward-looking statements. Risks and uncertainties that may cause actual results to differ materially include uncertainties inherent in the development of therapeutic product candidates, such as preclinical discovery and development, conduct of clinical trials and related regulatory requirements, our reliance on third parties over which we may not always have full control, and other risk and uncertainties that are described in our Annual Report on Form 10-K filed with the SEC on March 30, 2021 and our other Periodic Reports filed with the SEC. Forward-looking statements represent our management's beliefs and assumptions only as of the date of this presentation and include all matters that are not historical facts. Our actual future results may be materially different from what we expect. Except as required by law, we assume no obligation to update these forward-looking statements publicly, or to update the reasons actual results could differ materially from those anticipated in the forward-looking statements, even if new information becomes available in the future.


Certain information contained in this presentation relates to, or is based on, studies, publications, surveys and other data obtained from third-party sources and the Company's own internal estimates and research. While the Company believes these third-party sources to be reliable as of the date of this presentation, it has not independently verified, and makes no representation as to the adequacy, fairness, accuracy or completeness of, any information obtained from third-party sources. In addition, all of the market data included in this presentation involves a number of assumptions and limitations, and there can be no guarantee as to the accuracy or reliability of such assumptions. Finally, while we believe our own internal research is reliable, such research has not been verified by any independent source.

Two Platforms to Unleash Anti-Cancer T-cell Activity



TMAb™ (Tumor Microenvironment Activated Biologics) Platform

- Next-generation tumor activated mAbs
- Binding only in the low-pH tumor microenvironment
- Target checkpoints and/or other immune pathways
- Enable improved PK/PD and toxicity profiles




ImmunoPhage™ Platform

- Powerfully self-adjuvanted nanoparticle vaccine can drive B cell and T cell responses
- Multi-antigen vaccine enables personalized approach from “off-the-shelf” components
- Targets APCs
- Enhanced through addition of immunostimulatory nanobodies & cytokines


T-Cells Are Central to Our Approach and the Key to Unlocking Groundbreaking Clinical Activity

TMAb



Focus on novel immune checkpoints to
UNLEASH anti-tumor T-cells

ImmunoPhage™



Focus on multi-antigen approach for HLA-mediated immunotherapy to **GENERATE** anti-tumor T-cells

Positioned to Drive Value with Next Generation Product & Platform Development



Two Platforms Provide Broad Pipeline Opportunities

**TMAb*
Platform**

**ImmunoPhage™
Platform**

**In-house GMP
manufacturing
capabilities**



Strong Cash Position





Business Development

*Tumor Microenvironment Activated biologics

Pipeline Utilizing Pioneering ImmunoPhage Platform, TMAb Platform



	Program (Target)	Indication	Discovery	IND-enabling	Phase 1 / 2 Clinical
 TMAb	SNS-101 (VISTA)	Solid Tumors	<div></div>		
	SNS-VSIG4	Solid Tumors	<div></div>		
 ImmunoPhage	SNS-401-NG (Multiple Tumor Antigens)	Merkel Cell Carcinoma	<div></div>		
		Head and Neck Cancer	<div></div>		
		Lung Cancer	<div></div>		
		Melanoma	<div></div>		
		Breast Cancer	<div></div>		


TMAb (Tumor Microenvironment Activated biologics) Platform



pH-sensitive Antibodies Only Bind their Targets in the Low-pH Tumor Microenvironment

TMAb PLATFORM

The tumor microenvironment of pH 6.0 is lower than physiological pH of 7.4




Sensei's technology identifies pH-sensitive antibodies that bind only at the tumor

- Antibodies that bind at physiological pH may encounter a “sink”
 - Prevents effective binding at the tumor and may lead to toxicity
- TMAb antibodies bypass tissue compartments other than the low-pH tumor microenvironment
- Potential for improved safety and clinical activity profile

Why a pH-sensitive Antibody is Important

TMAb Platform


- Antibodies that bind at physiological pH may result in rapid elimination from circulation through targeted-mediated drug disposition (TMDD)
- In such cases, efficacious drug occupancy levels may be difficult to reach, potentially narrowing the therapeutic window



VISTA: An Emerging Checkpoint Target on Myeloid Cells

- First TMAb™ program against VISTA
- B7 family ligand
- Expressed on myeloid cells, macrophages, NK cells and T-regs¹
- Inhibition of VISTA may lead to activation of myeloid cells
- Excellent therapeutic combinability with CTLA-4 or PD-1/PD-L1 ICIs, especially in cold tumors²
- VISTA expression correlates with poor survival rates across multiple cancers
- **Novel development program with no approved therapies**

VISTA is a Negative Regulator of T cell Function



¹ Lines et al. Cancer research vol. 74,7 (2014)

² Gao et al. Nature medicine vol. 23,5 (2017)

Increased Understanding of VISTA as a Promising Target to Address the Needs of Patients with Cancer



nature
medicine

© 2017 Nature America, Inc., part of Springer Nature. All rights reserved.

nature
medicine

BRIEF COMMUNICATIONS

VISTA is an inhibitory immune checkpoint that is increased after ipilimumab therapy in patients with prostate cancer

Jianjun Gao¹, John F Ward², Curtis A Pettaway³, Lewis Z Shi¹, Sumit K Subudhi¹, Luis M Vence⁴, Hao Zhao⁵, Jianfeng Chen¹, Hong Chen⁶, Eloni Hristova⁶, Patricia Troncoso⁶, James P Allison^{1,3}, Christopher J Logothetis¹, Ignacio I Wistuba⁶, Manuel A Sepulveda², Jingjing Sun⁷, Jennifer Wargo⁸, Jorge Blando⁹ & Padmanee Sharma^{1,3,5}

To date, anti-CTLA-4 (ipilimumab) or anti-PD-1 (nivolumab) monotherapy has not been demonstrated to be of substantial clinical benefit in patients with prostate cancer. To identify additional immune-inhibitory pathways in the prostate-tumor microenvironment, we evaluated untreated and ipilimumab-treated tumors from patients in a presurgical clinical trial. Levels of the PD-L1 and VISTA inhibitory molecules increased on independent subsets of macrophages in treated tumors. Our data suggest that VISTA represents another compensatory inhibitory pathway in prostate tumors after ipilimumab therapy.

Immune checkpoint therapies, including anti-CTLA-4 and anti-PD-1 therapies, that block T cell inhibitory pathways have led to durable antitumor responses and clinical benefit in a substantial number of patients with cancer^{1,2}. However, prostate cancer has proven to be poorly responsive to immune checkpoint monotherapy^{3–5}. To better understand the immune profile within prostate tumors and potential compensatory immune-inhibitory pathways that may arise in the setting of immune checkpoint monotherapy, we conducted a clinical trial (NCT01194271) with ipilimumab plus androgen-deprivation therapy (ADT) before surgery in patients with localized prostate cancer (Supplementary Fig. 1a–c and Supplementary Tables 1 and 2).

We compared post-treatment and baseline blood samples (Supplementary Fig. 1a), evaluating the levels of CD4⁺ and CD8⁺ T cells (Supplementary Fig. 2a), as well as those of T cell subsets expressing inducible costimulator (ICOS), OX40, 4-1BB, PD-1, CTLA-4, and FoxP3 (Supplementary Fig. 2a,b). We observed an increase in CD4⁺ and CD8⁺ T cells, including PD-1⁺ and ICOS⁺ subsets, after ipilimumab therapy, which is similar to our previous findings with ipilimumab monotherapy in patients with melanoma

and bladder cancer^{6–8}. We also compared post-treatment tumor tissues (Supplementary Fig. 1a) to those of stage-matched untreated tumors from another cohort of patients (Supplementary Fig. 1b). Flow cytometric studies revealed a significantly higher frequency of CD4⁺, CD8⁺, and ICOS⁺ T cells in the post-treatment tumors (Fig. 1a). Immunohistochemical (IHC) studies also demonstrated significant increases in tumor-infiltrating immune cells, including CD4⁺, CD8⁺, ICOS⁺, CD45RO⁺, granzyme-B (GrB)⁺, and CD68⁺ cells (Supplementary Fig. 3). We found significantly greater immune cell infiltration in prostate tumors after ipilimumab therapy but not after ADT alone, although ADT monotherapy was associated with significantly higher levels of ICOS⁺ and GrB⁺ cells, which may represent an activated T cell subset (Fig. 1b). Taken together, our data suggest that the immunologic changes in post-treatment tumors were mostly due to ipilimumab therapy, as opposed to ADT. However, we cannot discount a possible synergistic effect between ipilimumab and ADT.

We did not observe clinical responses consisting of pathologic complete response, as we did previously for patients with bladder cancer⁸. To identify potential mechanisms that might explain this lack of response, we performed an unbiased gene expression study and found that ipilimumab therapy resulted in significant changes in the expression of a total of 690 genes (false discovery rate (FDR) <0.2; $P < 0.028$; log₂ (fold change) > 0.5) (Supplementary Table 3), most of which are related to immune responses (Supplementary Fig. 4a). We focused our analyses on a subset of genes that represent inhibitory immune checkpoints and identified increased PD-L1 and VISTA expression in post-treatment tumors (Supplementary Fig. 4b). Both PD-L1 and VISTA were previously reported as inhibitory molecules that can suppress murine and human T cell responses^{9,10}. Here we found significantly greater protein expression of PD-1, PD-L1, and VISTA in prostate tumors after ipilimumab therapy (Fig. 1c and Supplementary Fig. 5a).

We also evaluated metastatic tumors and blood samples from patients with metastatic prostate cancer who took part in a separate clinical trial (NCT02113657) and received treatment with ipilimumab, finding an increase in PD-L1 and VISTA expression in tumor tissues (Supplementary Fig. 5b) as well as on monocytes in blood (Supplementary Fig. 6a), which was similar to data from a mouse model of prostate cancer (Supplementary Fig. 6b). We suggest that PD-L1 and VISTA are likely to be relevant inhibitory immune checkpoints in both localized and metastatic prostate cancer.

We evaluated PD-L1 and VISTA expression in different cell subtypes from matched pre- and post-treatment prostate tumors and observed significantly higher PD-L1 expression on CD4⁺ T cells, CD8⁺ T cells, and CD68⁺ macrophages after treatment (Supplementary Fig. 7a).

¹Department of Genitourinary Medical Oncology, University of Texas MD Anderson Cancer Center, Houston, Texas, USA. ²Department of Urology, University of Texas MD Anderson Cancer Center, Houston, Texas, USA. ³The Immunotherapy Platform, University of Texas MD Anderson Cancer Center, Houston, Texas, USA. ⁴Department of Pathology, University of Texas MD Anderson Cancer Center, Houston, Texas, USA. ⁵Department of Immunology, University of Texas MD Anderson Cancer Center, Houston, Texas, USA. ⁶Department of Translational Molecular Pathology, University of Texas MD Anderson Cancer Center, Houston, Texas, USA. ⁷Janssen Oncology Therapeutic Area, Janssen Research and Development, LLC, Pharmaceutical Companies of Johnson & Johnson, Spring House, Pennsylvania, USA. ⁸Department of Surgical Oncology, University of Texas MD Anderson Cancer Center, Houston, Texas, USA. Correspondence should be addressed to P.S. (ps@mdanderson.org).

Received 16 December 2016; accepted 17 February 2017; published online 27 March 2017; doi:10.1038/nm.4308

Trends in
Immunology

CellPress
OPEN ACCESS

Feature Review

VISTA: A Mediator of Quiescence and a Promising Target in Cancer Immunotherapy

Long Yuan,^{1,2} Jahnvi Tatineni,² Kathleen M. Mahoney,^{2,3} and Gordon J. Freeman^{2,4*}

V-domain Ig suppressor of T cell activation (VISTA) is a B7 family member that maintains T cell and myeloid quiescence and is a promising target for combination cancer immunotherapy. During inflammatory challenges, VISTA activity reprograms macrophages towards reduced production of proinflammatory cytokines and increased production of interleukin (IL)-10 and other anti-inflammatory mediators. The interaction of VISTA with its ligands is regulated by pH, and the acidic pH ~6.0 in the tumor microenvironment (TME) facilitates VISTA binding to P-selectin glycoprotein ligand 1 (PSGL-1). Targeting intratumoral pH might be a way to reduce the immunoinhibitory activity of the VISTA pathway and enhance antitumor immune responses. We review differences among VISTA therapeutics under development as candidate immunotherapies, focusing on VISTA binding partners and the unique structural features of this interaction.

VISTA: How This B7 Protein Might Transform Cancer Immunotherapy

Immunotherapy has become an established pillar of cancer treatment, in large part owing to the success of blocking the programmed cell death protein 1 (PD-1)/programmed death-ligand 1 (PD-L1) immune checkpoint (see Glossary) pathway. As recent research deepens our understanding of V-domain Ig suppressor of T cell activation (VISTA), the VISTA signaling pathway has increasingly become a promising target for overcoming resistance to current immune checkpoint therapies [1]. Although the development of VISTA blocking antibodies has not reached fruition clinically, this review highlights the new features of VISTA that make this pathway particularly attractive for therapeutic development. We discuss (i) VISTA expression on immune cells in the tumor microenvironment (TME), (ii) the biological functions and bidirectional signaling pathways of VISTA in mammalian lymphocytes and myeloid cells, (iii) the structural features of VISTA that contribute to its molecular interactions, (iv) current VISTA monoclonal antibodies (mAbs) that are in clinical development, and (v) the candidate druggable targets that regulate the pH of the TME and which in turn might affect VISTA activity in vivo. This review gives a detailed picture of VISTA structure in the context of its binding partners and therapeutic antibodies targeting VISTA.

VISTA Structure

VISTA, also known as PD-1H, B7-H5, Diest1, G24, DD10, and C10orf54, is encoded by the VSIF gene in human (Vslr in mouse) and has multiple unique features, including its interaction with two receptors that bind to overlapping but distinct sites on the VISTA extracellular domain (ECD) [2–4]. VISTA is a type I transmembrane protein that was identified by mRNA analysis of activated versus resting mouse natural regulatory T cells (Tregs) [5] and also by homology to coinhibitory molecules such as PD-1 [6]. VISTA bears features of both the B7 and CD28 families of immunoregulatory molecules and can act as both a ligand and a receptor [3, 7, 8]. The VISTA ECD is most homologous to the B7 family, which includes well-known immune checkpoint ligands such as PD-L1 (Figure 1c). Whereas other B7 family members have an IgV-like and IgC-like domain, mouse and human VISTA contain a single unusually large IgV-like domain (Figure 1a) [2]. VISTA

Highlights

V-domain Ig suppressor of T cell activation (VISTA) binds to V-set and Ig domain-containing 3 (VSG3) and P-selectin glycoprotein ligand 1 (PSGL-1) ligands, and signaling may be bidirectional.

VISTA binds to PSGL-1 at acidic pH, such as in the tumor microenvironment (TME), but not at physiological pH.

VISTA activity imposes quiescence on mammalian myeloid and naive T cells, and inhibits T cell activation and cytokine production. It can promote peripheral tolerance via enhanced activation-induced T cell death.

VISTA is particularly upregulated on myeloid-derived suppressor cells (MDSCs) via hypoxia, and can contribute to the immunosuppressive functions of myeloid cells by reducing Toll-like receptor (TLR) signaling and cell migration, as well as by reprogramming myeloid cells towards reduced production of the proinflammatory cytokines interleukin (IL)-6, tumor necrosis factor (TNF)- α , and IL-12, and increased production of IL-10 and other anti-inflammatory mediators.

Antagonistic VISTA antibodies are in clinical development for treating some cancers; drugs that target the acidity of the TME might reduce immunoinhibitory activity in acidic niches and combine well with VISTA or checkpoint blockade therapies.


¹Program in Immunology, Harvard Medical School, Boston, MA 02115, USA

²Department of Medical Oncology, Dana-Farber Cancer Institute, Harvard Medical School, Boston, MA 02215, USA



VISTA Checkpoint is Activated at the Low pH of the Tumor Microenvironment

Antibodies that block VISTA histidines:
H153, H154 and H155 on interrupt
PSGL-1 binding¹




VISTA's extracellular domain is uniquely rich in histidines¹

Histidines are protonated at low pH enabling VISTA to distinguish the active (acidic pH) and inactive (neutral pH) PSGL-1 binding interface

Anti-VISTA Programs in Development




TMAb Platform

		BMS	Kineta	CI-8993 (formerly JNJ-61610588)	PF-W0180	HMBD-002
pH Sensitivity	Yes	Yes	No	No	No	No
Stage	Preclinical	Preclinical	Preclinical	Phase I	Phase I	Phase I
Clinical Data / Notes	<ul style="list-style-type: none"> Preclinical data to be presented by year-end 2021 IND-enabling studies to initiate by year-end 2021 	<ul style="list-style-type: none"> N/A 	<ul style="list-style-type: none"> N/A 	<ul style="list-style-type: none"> JNJ initiated Phase I study in 2016 12 pts enrolled; initial dose was 0.005 mg/kg Only patient treated at 0.3 mg/kg experienced grade 3 CRS-associated encephalopathy and trial was halted 	<ul style="list-style-type: none"> Ongoing; no data reported 	<ul style="list-style-type: none"> Ongoing; no data reported

pH-Sensitive Anti-VISTA Antibodies Showed Positive Results *In Vivo*




Sensei Anti-VISTA Parental mAb Tumor Growth of MC38 in Hu VISTA Knock-in Mice




Dosing: 14d @ 20mpk q3d

Parental mAb is pH-Sensitive

Antigen binding at pH 6.0




Antigen binding at pH 7.4



Significant binding occurred at low pH within the tumor microenvironment; No significant binding observed at physiological pH (7.4)

VSIG4: A Novel Next Generation Checkpoint Modulating the Tumor Microenvironment




No approved therapies against VSIG4

- Second TMAb program
- B7 family related protein
- Expressed on macrophages
- Inhibits T-cell activation
- Novel therapeutic combinability with existing IO drugs

ImmunoPhage™ Platform



Bacteriophage




Ubiquitous viruses that infect bacteria but not mammalian cells. Adept at activating the human immune system in multiple unique ways

Generating Strong Antibody and T-cell Responses

ImmunoPhage Platform


Bacteriophage virus is engineered and manufactured with both antigen and immune stimulatory viral DNA



The **ImmunoPhage™** bacteriophage is an icosahedron with a tail. This configuration can be viewed as an activating signal to the immune system

Generating Strong Antibody and T-cell Responses

ImmunoPhages are taken-up by APCs and deliver three critical signals required to drive activation of T cells.



ImmunoPhage™ A Multi-Pronged Approach to Address the Complexities of Cancer



Our **ImmunoPhages** can mount a multi-modal attack on cancer, combining the benefits of a traditional vaccine with localized gene therapy

Targeted therapeutic vaccine

- MHC-mediated immunity
- Bacteriophage have natural tropism for APCs
- Can be further targeted to APCs with non-antigen capsid modifications



Phortress™ library

- Personalized - yet off the shelf - medicines
- Pre-manufactured cost effectively - then combined based on genetic profile

Gene therapy vehicle

- Phage containing self-replicating RNA
- Used to deliver payloads consisting of immunomodulatory proteins or nanobodies

SNS-401-NG: Building the First Custom Merkel Cell Polyoma Virus (MCPyV) ImmunoPhage



SNS-401-NG Development



Collaboration with University of Washington to build **first custom Merkel Cell Carcinoma (MCC) vaccine consisting of Merkel Cell Polyoma Virus epitopes** and other patient specific antigens

MCC is a rare, aggressive neuroendocrine skin cancer

- 33-46% disease-specific mortality
- 2,500 cases/yr with disease-specific mortality approaching 50%
- Vaccine combination therapy in adjuvant or neoadjuvant is attractive and feasible
 - PD-1/PD-L1 refractory MCC remains unmet medical need with aggressive clinical course
 - ~40% MCC patients recur <24 months following definitive local treatment




Integration of MCPyV is present in ~80% of U.S. cases

- In these cases, expression of a viral antigen (oncogenic T-antigen) **appears to be a strictly required tumor driver**
- Researchers at UW have mapped MCPyV epitopes and **determined CD8 T-cell, CD4 T-cell, and B-cell epitopes that are antigenic** in the context of MCPyV+ MCC tumors.





Mechanism of Action: Activation and Maturation of Dendritic Cells

Dose-response of engineered lambda phage on human skin-derived DC cultures

Signal: Dendritic cell co-stimulatory molecules



Signal: Cytokine secretion




Critical signals of dendritic cell activation show dose-dependent increases when cells are exposed to increasing amounts of ImmunoPhages

SNS-401-NG has Potential to be First Fully Customized, Yet Off-the-Shelf, Therapy




SNS-401-NG Development in Merkle Cell

Patients would receive a bespoke mixture of ImmunoPhage that included antigens from the MCPyV and a subset of TAA-expressing ImmunoPhage




Most MCC tumors contain multipleTAAs¹



1. Based on internal data

Phortress: Proprietary Library of Personalized Vaccine Cocktails with Off-the-Shelf ImmunoPhage “ingredients”



- These “cocktails” are defined by the disease or patient genetics
- Combinations are customized to cover multiple epitopes, protein domains or targets
- Each *ImmunoPhage* is pre-manufactured to target a discrete antigen

Personalized Immunotherapy Approach Could Accelerate Speed to Treatment




High speed and low cost-of-goods of ImmunoPhage allows a broader array of antigens

Personalized yet Off-the Shelf TAA Therapy

Off-the-Shelf + Patient-specific Neoantigen Therapy

Routine Biopsy




Clinical biopsy of tumor
as input material

Tumor Sequencing




Tumor DNA
Tumor RNA
Normal DNA

Personalized yet Off-the-shelf ImmunoPhage Cocktail




Assemble a
personalized cocktail
from off-the-shelf TAA
ImmunoPhage for
administration

Neoantigen Prediction



Identify additional tumor
specific neoantigens

Neoantigen ImmunoPhage Manufacturing




Engineer novel
ImmunoPhages
expressing distinct tumor
specific epitopes

ImmunoPhage Injection Including Neoantigens



Deliver neoantigen
ImmunoPhage cocktail
for administration and
add neoantigen phages
to bank for future use

Sensei's Vision to Capture Platform and Pipeline Value



Proven Team With Deep Experience



John Celebi, MBA
President and CEO



Michael Boychyn, PhD
SVP, CMC



Erin Colgan
SVP, Finance and Administration



Pauline Callinan, PhD
VP, Business Operations and Strategy



Marie-Louise Fjaellskog, MD, PhD
Chief Medical Officer



Jean Campbell, PhD
VP, Biologics Discovery



Elisabeth Colunio
VP, Human Resources



Alice Drumheller
VP, Clinical Operations



Robert Pierce, MD
Chief Scientific Officer



Bao Le
VP, Regulatory



Lora Pike
VP, Investor Relations Communications



Edward van der Horst, PhD
VP, Preclinical Development

Upcoming Expected Program Milestones



SNS-101 (anti-VISTA)

YE 2021:

- Present preclinical data at scientific meeting
- ✓ Select lead candidate
- Initiate IND-enabling studies



SNS-401-NG

2H 2022:

- initiate IND-enabling studies



SNS-VSIG4

2023:

- Select product candidate



Training the Immune System to Fight Cancer

John K. Celebi, MBA
President & Chief Executive Officer

September 2021

NASDAQ: SNSE

